AD		

Award Number: DAMD17-02-1-0245

TITLE: A Phase II Immunotherapeutic Trial: Combination Androgen

Ablative Therapy and CTLA-4 Blockade as a Treatment for

Advanced Prostate Cancer

PRINCIPAL INVESTIGATOR: Eugene D. Kwon, M.D., Ph.D.

CONTRACTING ORGANIZATION: Mayo Clinic Rochester

Rochester, MN 55905

REPORT DATE: December 2003

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20040901 065

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY	2. REPORT DATE	3. REPORT TYPE AND	DATES COVERE	
(Leave blank)				
	December 2003	Annual (1 Dec	2002 - 30 N	lov 2003)
4. TITLE AND SUBTITLE			5. FUNDING N	UMBERS
A Phase II Immunotherap	eutic Trial: Combina	tion Androgen		
Ablative Therapy and CT	TLA-4 Blockade as a '	Treatment for	DAMD17-02-1-0245	
Advanced Prostate Cancer				
6. AUTHOR(S)			1	
				•
Eugene D. Kwon, M.D., Ph	1.D.			
, ,				
7. PERFORMING ORGANIZATION NAM	ME(S) AND ADDRESS(ES)		8. PERFORMING	G ORGANIZATION
Mayo Clinic Rochester	•••		REPORT NUI	
Rochester, MN 55905				·
1.00				
E-Mail: kwon.eugene@mayo.	. edu			
9. SPONSORING / MONITORING	2/50)		10. SPONSORING / MONITORING	
AGENCY NAME(S) AND ADDRESS	• •		AGENCY R	EPORT NUMBER
U.S. Army Medical Resear		and		
Fort Detrick, Maryland	21702-5012			
11. SUPPLEMENTARY NOTES				•
12a. DISTRIBUTION / AVAILABILITY	STATEMENT			12b. DISTRIBUTION CODE
Approved for Public Release; Distribution Unlimited				
	•			
				<u> </u>
13. ABSTRACT (Maximum 200 Word	s)			
The induction of tumor-	-specific T cells rema	ins a primary c	bstacle to	immunotherapeutic
approaches for most cancers including prostate cancer. This difficulty has been largely				
ascribed to mechanisms for tumor evasion of the immune system and host-imposed				
restrictions (collectively referred to as tolerance) that prevent cross-reactive				
autoimmunity against the parent tissues from which tumors arise. Limitations in				
techniques to identify novel and truly immunogenic prostate-specific antigens and				
efficient methods to modify autologous tissues for vaccine preparation have further				

approaches for most cancers including prostate cancer. This difficulty has been largely ascribed to mechanisms for tumor evasion of the immune system and host-imposed restrictions (collectively referred to as tolerance) that prevent cross-reactive autoimmunity against the parent tissues from which tumors arise. Limitations in techniques to identify novel and truly immunogenic prostate-specific antigens and efficient methods to modify autologous tissues for vaccine preparation have further constrained approaches to develop immune-based therapies for prostate cancer. Hence, relatively straightforward manipulations that induce specific T cell responses against prostate tumors or epithelial tissues, especially in vivo, might ultimately prove valuable for prostate cancer immunotherapy. This study explores combined androgen ablation (AA) + CTLA-4 blockade immunotherapy as a means of potentiating T cell-mediated responses against prostate tumors to improve overall treatment of advanced prostate cancer.

14. SUBJECT TERMS Prostate Cancer			15. NUMBER OF PAGES 8
Flostate Cancel			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT
Unclassified	Unclassified	Unclassified	Unlimited

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89) Prescribed by ANSI Std. Z39-18 298-102

Table of Contents

Gover1
SF 2982
Table of Contents3
Introduction4
Body5
Key Research Accomplishments7
Reportable Outcomes7
Conclusions7
References7
Appendicesnone

INTRODUCTION.

Although androgen ablation (AA) has long served as a mainstay of treatment for patients with advanced prostate cancer, this form of therapy is only palliative. Consequently, nearly all of the ~35,000 men who will die of prostate cancer each year will do so after failing AA therapy. For patients with advanced prostate cancer, median duration of treatment response averages ~2 years following institution of AA therapy. Upon failure of AA therapy, when metastatic hormone-refractory prostate cancer progression becomes apparent, median survival averages only another 16-20 months. While a number of investigational agents have demonstrated some promise for patients who have failed AA, to date, no agent or combined regimen has been shown to confer a survival advantage in randomized clinical testing. Thus, novel approaches for treating advanced prostate cancer — particularly interventions capable of acting synergistically with AA therapy -- are needed. This clinical phase II trial will test whether a specific form of immunotherapy, CTLA-4 blockade, can be used to enhance treatment responses in patients with advanced prostate cancer being treated with AA therapy. CTLA-4 is a T-cell receptor that mediates downregulation of T-cell function. CTLA-4 blockade refers to the administration of a monoclonal antibody (MDX- CTLA-4) that binds the T-cell CTLA-4 receptor to prevent inhibition of T-cell function.

It has been established that CTLA-4 blockade immunotherapy is capable of potentiating host T cell-mediated responses against a number of tumors including prostate cancer, and that the antitumoral capability of CTLA-4 blockade can be markedly enhanced by combining CTLA-4 blockade with manipulations that induce tumor/tissue-specific T cells. Finally, it has recently been shown that androgen ablative (AA) therapy promotes prostate tissue infiltration by T cells that may exhibit specific activity against prostate epithelial/tumor cells. It is, therefore, hypothesized that AA therapy can be used to coax prostate tissues to behave as "in situ vaccines" to induce prostate tissue/tumor-specific T cells (refer to schematic; right). Such prostate-specific T cells induced by AA therapy might then be amenable to immunotherapeutic potentiation by CTLA-4 blockade. If so, T cell responses raised by AA therapy and then bolstered by CTLA-4 blockade would be expected to contribute to tumor rejection, ultimately manifesting in improved treatment outcome for patients with advanced prostate cancer.

In contrast, T cell responses in patients treated with AA therapy alone would be expected to wane, ultimately giving way to tumor progression. Hence, to test whether combined AA therapy plus CTLA-4 blockade is capable of eliciting clinical treatment responses superior to those provided by standard AA therapy alone, and to further test that such treatment advantages emanate from potentiation of host T cell-mediated antitumoral immunity by CTLA-4 blockade, the following prospective randomized clinical phase II trial is proposed.

The primary endpoint for this study is proportion of patients remaining free of disease progression. The time to disease progression (progression-free interval), is defined as the interval from the start of AA therapy until PSA progression. Secondary endpoints will include assessments of initial PSA response to treatment. Tertiary endpoints will include exploratory correlative assessments of prostate-specific as well as generalized host T cell response to treatment. For the cross over component of this trial in which control group subjects will receive CTLA-4 blockade upon progression, the cross over study endpoint will be an alternate definition of PSA response (regression, stabilization or progression). The immunologic studies proposed in this protocol may prove relevant to recent observations suggesting that clinical responses to AA therapy are, at least in part, predicated upon an immune response by the host. If true, immunologic endpoints in this study might ultimately be applicable for identifying patients at greatest risk of failing AA therapy, and who might benefit from early and aggressive intervention.

BODY.

We are currently prepared to open accrual onto our phase II trial entitled "A Phase II Immunotherapeutic Trial: Combination Androgen Ablative Therapy and CTLA-4 Blockade as a Treatment for Advanced Prostate Cancer" here at Mayo. It is also anticipated that accrual onto this study will open shortly at UCSF (E. Small, PI) which represents the sister site for this protocol. Until now, our efforts have largely been dedicated towards final development of our phase II protocol, as well as acquistion of requisite approval from the respective IRB's at Mayo, UCSF and the DOD. Additionally, we have requested and secured an IND from the FDA that authorizes our study of combined MDX-CTLA-4 and AA therapy for treatment of advanced prostate cancer. The key steps that we have dealt with in order to open this study are outline in what follows below.

•	01/18/01	Concept approved by the Cancer Center Support Grant Concept Review Committee
•	07/31/02	Submitted for review by the Mayo Cancer Center Protocol Review Committee
•	08/07/02	Approved by the Mayo Cancer Center Protocol Review Committee with minor modifications and pending sign-off by Dr. J. O'Fallon
•	08/23/02	Protocol submitted for IRB approval
•	08/23/02	IRB deferred; further consideration will require 13 items (see minute/file)
•	08/25/02	Dr. Kwon application for IND (cross-referencing Medarex BB-IND 8937).
•	08/29/02	IRB approval received; correspondence involved 10 items; CF approved with revisions
•	09/10/02	Assignment of IND BB-IND 10,655 to Dr. Kwon.
•	10/03/02	Communication with FDA (Dr. Lee Pai Scherf, Medical reviewer for IND) to clarify protocol.
•	10/05/02	Clarification of protocol and changes submitted to FDA (Dr. Pai Scherf).

• 10/31/	IRB approved Investigator made	Brochure (Edition 3.0); no CF revisions were
• 11/07/	by the Human Subjects Res	litional information prior to scheduling for review earch Review Board (HSRRB). Issue pertaining ted injury raised by Mayo and DOD.
• 03/06/	IRB approved consent form	changes to comply with HIPAA
• 04/17/	Revised clinical protocol, re sent to Andrea Kline	evised consent, and responses to the DOD IRB
• 05/05/		dditional information prior to Human Subjects SRRB) meeting scheduled for 05/14/03
• 05/05/	Dr. Kwon provided additiona	l information to Andrea Kline
• 05/09/	Andrea Kline forwarded tw review of the protocol	o questions from an MSRRB member's pre-
• 05/12/	Dr. Kwon responded to And IRB pre-review questions	drea Kline and provided a response to the DOD
• 05/14/	Dr. Kwon teleconference wi approval of protocol.	th the DOD HSRRB (IRB) board for review and
• 05/15/	Further clarifications of the to teleconference of 15/14/0	protocol submitted to DOD HSRRB in response 3.
• 06/12/	HSRRB approved revised p	rotocol
• 07/03/	versions to the Mayo Clinic the HSRRB requested. documentation of the IRB's consent form, the project ca	
*	office."	
• 07/21		
• 08/14/		
• 08/14/	OB IRB approval received for m	odification request dated 8/4/03 (see minute/file)
• 08/26	D3 BB-IND 10,655: Annual Re (Form FDA 1571).	eport of Progress (Serial Number 002) submitted
• 09/02	Dr. Kwon informed Andromodification	ea Kline that the Mayo IRB approved the
• 09/19	Dr. Kwon forwarded the Ma Kline	ayo IRB approval minute and consent to Andrea
• 09/25	O3 Andrea Kline questioned r consent	missing HSRRB-recommended changes to the
• 09/25	Dr. Kwon responded that s the consent were unaccepta	some of the HSRRB-recommended changes to able to the Mayo IRB
• 09/30	O3 Mr. Tibor Tuzson asked if a the Mayo IRB; there were d	in older version of the consent was approved by iscrepancies
• 10/09	Modification received by the	RB
• 10/30	· · · · · · · · · · · · · · · · · · ·	modification dated 10/6/03; involves revisions to
• 11/11	O3 IRB approval minutes forwa	rded to Andrea Kline
• 11/18	Dr. Kwon received an e-m	nail from Andrea Kline stating that an approval sent to the USAMRAA Contracting Office and

that this office would contact the Mayo Clinic grant office regarding permission to implement this protocol in the near future.

02/13/04

Sharri Hackbarth, Mayo Research Administrator, received letter from USAMRAA with "copies of Modification No. P00001 for Grant No DAMD17-02-1-0245 fully executed for your information and file."

KEY RESEARCH ACCOMPLISHMENTS.

- Protocol written and fully approved by the Mayo IRB and DOD HSRRB.
- Resolved issues pertaining to patient coverage for research-related injury.
- Secured IND BB 10,655 from FDA.
- Protocol in final stage of approval by UCSF IRB.

REPORTABLE OUTCOMES TO DATE.

None

CONCLUSIONS.

In summary, we have now accomplished the necessary steps to open our phase II trial entitled "A Phase II Immunotherapeutic Trial: Combination Androgen Ablative Therapy and CTLA-4 Blockade as a Treatment for Advanced Prostate Cancer" here at Mayo. Accrual onto this study will also open at UCSF very shortly. It is our sincere hope that this trial will ultimately demonstrate that T cell-mediated responses against prostate tumors, induced by an inductive period of androgen ablation, will be immunotherapeutically potentiated by anti-CTLA-4 mAb therapy to improve overall treatment for patients that initially present with advanced prostate cancer.

RELEVANT REFERENCES.

Parker SL, Tong T, Bolden S, and Wingo PA. Cancer Statistics, 1997. CA Cancer J. Clin., 47:5-27, 2000.

McLeod G and Obrien-Abt ME. Hormonal management of metastatic prostate cancer and quality of life issues. Comprehensive Textbook of Genitourinary Oncology. Vogelzang NJ, Scardino PT, Shipley WU, Coffey DS, eds. Williams & Wilkins, Baltimore MD. pp854-874, 1996.

Dawson NA and Vogelzang NJ. Prostate Cancer. Wiley-Liss, New York, 1999.

Oliver RTD and Gallagher CJG. Intermittent endocrine therapy and its potential for chemoprevention of prostate cancer. Canc. Surv., 23:191-207, 1995.

Wise GJ, Marella V.K, Talluri G and Shirazian D. Cytokine variations in patients with hormone treated prostate cancer. J. Urol. 164:722-725, 2000.

Vesalainen S, Lipponen P, Talja M and Syrjanen K. Histological grade, perineural infiltration, tumour-infiltrating lymphocytes and apoptosis as determinants of long-term prognosis in prostatic adenocarcinoma. Eur. J. Cancer 30A:1797-1803, 1994.

Shimura S, Yang G, Ebara S, Wheeler TM, Frolov A and Thompson TC. Reduced infiltration of tumor-associated macrophages in human prostate cancer: Association with cancer progression. Cancer Res. 60(20):5857-61, 2000.

Kwon ED, Foster BA, Hurwitz AA, Madias C, Allison JP, Greenberg NM, Burg MB. Elimination of residual metastatic prostate cancer following surgery and adjunctive CTLA-4 blockade immunotherapy. Proc. Natl. Acad. Sci. USA 96:15074, 1999.

Hurwitz AA, Yu TF, Leach DR, Allison. CTLA-4 blockade synergizes with tumor-derived GM-CSF for treatment of an experimental mammary carcinoma. Proc. Natl. Acad. Sci. USA 95:10067-10071, 1998.

van Elsas A, Hurwitz AA, Allison JP. Combination immunotherapy of B 16 melanoma using anti-CTLA-4 and GM-CSF producing vaccines induces rejection of subcutaneous and metastatic tumors accompanied by autoimmune depigmentation. J. Exp. Med. 190:355-366, 1999.

Hurwitz AA, Foster BA, Kwon ED, Trong T, Choi EM, Greenberg NM, Burg MB, Allison JR. Combination immunotherapy of primary prostate cancer in a transgenic model using CTLA-4 blockade. Cancer Res. 60:2444-2448, 2000.

Hurwitz AA, Foster BA, Allison JP, Greenberg NM and Kwon ED. Use of the transgenic adenocarcinoma of mouse (TRAMP) model and its derivative for testing immunotherapies against prostate cancer. Current Protocols in Immunology. Current Protocols and John Wiley & Sons, Editor A.M. Kruisbeek. In Press, 2001.

Foster BA, Gingrich JR, Kwon ED, Madias C and Greenberg NM. Characterization of prostatic epithelial cell lines derived from transgenic adenocarcinoma of the mouse prostate (TRAMP) model. Cancer Res. 57:3325-3330, 1997.

Hurwitz AA, Kwon ED and Elsas AV. Costimulatory wars: the tumor menace. Current Opinion in Immunology. 12:589-596, 2000.

Mercader M, Bodner BK, Moser MT, Kwon PS, Park ES-Y, Maneke RG, Ellis TM, Wojcik EM, Yang D, Flanigan RC, Waters WB, Kast WM and Kwon ED. T cell infiltration of the prostate induced by androgen withdrawal in patients with prostate cancer. Proc. Natl. A cad. Sci. 98:14565-14570, 2001